

AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of the Claims

1. (Original) A method for stimulating perfusion, comprising the steps of:
selecting a patient having a tissue with compromised microvascular perfusion;
applying an ultrasound transducer to a location near the tissue; and
activating the transducer to initiate exposure of the tissue to ultrasound at a frequency
of 100 KHz to 2.0 MHz for a duration of 0.5 to 15 minutes, wherein local vasodilatation is
stimulated.
- 2-9. (Cancelled)
10. (Original) A method for stimulating reperfusion in a patient having a tissue
experiencing an ischemic injury, comprising the steps of:
selecting a patient experiencing a myocardial infarction;
applying a portable transcutaneous ultrasound transducer within 30 minutes after the
ischemic injury to a location near the tissue at the site of the ischemic injury;
activating the transducer to initiate exposure of the tissue to ultrasound and thereby
stimulate local vasodilatation; and
confirming the establishment of reperfusion.

11. (Original) The method of claim 10, further comprising the steps of applying a gel to enhance the transmission of ultrasound waves.

12. (Original) The method of claim 10, further comprising the steps of adjusting the focal length of the ultrasound waves.

13. (Original) The method of claim 10, wherein the frequency of the ultrasound waves is 100 KHz to 2.0 MHz.

14. (Original) The method of claim 10, wherein the ultrasound exposure is maintained for 15 minutes.

15. (Original) The method of claim 10, wherein the step of confirming the establishment of reperfusion comprises a procedure selected from the group consisting of angiography, electrocardiogram, diagnostic ultrasound, and measuring blood levels of creatine kinase.

16. (Original) The method of claim 10, further comprising the steps of injecting an anticlotting agent into the patient.

17. (Original) The method of claim 16, wherein the anticlotting agent is selected from the group consisting of aspirin, tissue plasminogen activator, and streptokinase.

18. (Original) A method for stimulating myocardial perfusion, comprising the steps of:

selecting a patient having a myocardium with compromised perfusion;
applying an ultrasound transducer to a location near the myocardium; and

activating the transducer to initiate exposure of the myocardium to ultrasound at a frequency of 100 KHz to 2.5 MHz, wherein myocardial blood flow is enhanced.

19. (Original) The method of claim 18, further comprising the step of confirming enhancement in myocardial blood flow.

20. (Original) The method of claim 18, wherein the transducer is activated to initiate exposure to ultrasound at a frequency of 100 KHz to 1.0 MHz.

21. (Original) The method of claim 18, wherein the transducer is activated to initiate exposure to ultrasound at a frequency of 100 KHz to 200 KHz.

22. (Original) The method of claim 18, wherein the exposure to ultrasound is for a duration of exposure is 0.5 to 15 minutes.

23. (Original) The method of claim 18, wherein the exposure to ultrasound is for a duration of exposure is 0.5 to 10 minutes.

24. (Original) The method of claim 18, wherein the exposure to ultrasound is for a duration of exposure is 5 to 10 minutes.

25. (Original) The method of claim 18, wherein the transducer is activated to initiate exposure to ultrasound with a temporal and spatial average energy level of 0.01 to 1.00 watts/cm².

26. (Original) The method of claim 18, wherein the transducer is activated to initiate exposure to ultrasound with pulsed modulation.

27. (Cancelled)

28. (Original) A method for stimulating cerebral perfusion, comprising the steps of:
selecting a patient having a cerebral tissue with compromised perfusion;
applying an ultrasound transducer to a location near the head; and
activating the transducer to initiate exposure of the head to ultrasound at a frequency
of 100 KHz to 2.5 MHz, wherein cerebral blood flow is enhanced.

29-37. (Cancelled)

38. (Original) A method for stimulating perfusion in a transplanted tissue,
comprising the steps of:
selecting a patient having a transplanted tissue;
applying an ultrasound transducer to a location near the transplanted tissue; and
activating the transducer to initiate exposure of the transplanted tissue to ultrasound at
a frequency of 100 KHz to 2.0 MHz for a duration of 0.5 to 15 minutes, wherein local vasodilatation
is stimulated.

39-51. (Cancelled)